

REPORT DOCUMENTATION PAGEForm Approved
OMB No. 0704-0188

The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to the Department of Defense, Executive Service Directorate (0704-0188). Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ORGANIZATION.

1. REPORT DATE (DD-MM-YYYY) 09/16/2017		2. REPORT TYPE Poster		3. DATES COVERED (From - To)	
4. TITLE AND SUBTITLE Pediatric Rapid Response Team: Vital Sign Based System vs. Pediatric Early Warning Score System				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
				5d. PROJECT NUMBER	
6. AUTHOR(S) Capt Scott Penney				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) 59th Clinical Research Division 1100 Willford Hall Loop, Bldg 4430 JBASA-Lackland, TX 78236-9908 210-292-7141				8. PERFORMING ORGANIZATION REPORT NUMBER 17310	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) 59th Clinical Research Division 1100 Willford Hall Loop, Bldg 4430 JBASA-Lackland, TX 78236-9908 210-292-7141				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release. Distribution is unlimited.					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON Clarice Longoria
a. REPORT	b. ABSTRACT	c. THIS PAGE			19b. TELEPHONE NUMBER (Include area code) 210-292-7141

MD; Mrs. Scarlett O'Hara-Wood, RN, BSN; Maj Lisa M. McFarlan, RN, BSN, MSNBC; Capt Robert Slaughter RN, BSN, CCRN;
Maj Carla S. Cox, RN, MS, PCNS-BC; 2Lt Amber N. Auge, RN, BSN; LtCol(S) Renée I. Matos, MD, MPH
Department of Pediatrics, Brooke Army Medical Center, Ft Sam Houston TX



- Approximately 8.5-14% of cardiopulmonary arrests in pediatrics occur outside the ICU with associated mortality rates from 50-67%*
- Only 10% of pediatric patients who suffer a cardiopulmonary arrest survive beyond one year post-event and 35% of survivors have neurological deficits*
- Pediatric rapid response teams (PRRTs) have been ineffective in preventing codes which decrease mortality in pediatric patients by 18%*
- The PRRT system was triggered by one abnormal vital sign (VS) parameter
- That limited nursing staff autonomy and critical thinking skills, resulting in the ineffective use of resources and staff
- Pediatric physiology easily prompts VS changes due to anxiety, fever, or medication delivery thus resulting in unnecessary PRRT activations
- Pediatric Early Warning Score (PEWS) system is an evidence-based tool shown to identify trends in patient hours preceding a cardiopulmonary event enabling earlier interventions* and prevention of further deterioration

- Approximately 8.5-14% of cardiopulmonary arrests in pediatrics occur outside the ICU with associated mortality rates from 50-67%*
- Only 10% of pediatric patients who suffer a cardiopulmonary arrest survive beyond one year post-event and 35% of survivors have neurological deficits*
- Pediatric rapid response teams (PRRTs) have been ineffective in preventing codes which decrease mortality in pediatric patients by 18%*
- The PRRT system was triggered by one abnormal vital sign (VS) parameter
- That limited nursing staff autonomy and critical thinking skills, resulting in the ineffective use of resources and staff
- Pediatric physiology easily prompts VS changes due to anxiety, fever, or medication delivery thus resulting in unnecessary PRRT activations
- Pediatric Early Warning Score (PEWS) system is an evidence-based tool shown to identify trends in patient hours preceding a cardiopulmonary event enabling earlier interventions* and prevention of further deterioration

- **Goal:** Implement a quality improvement initiative using the evidence-based PEWS criteria to improve recognition of deteriorating pediatric patients, allocation of PRRT resources, and pediatric staff satisfaction regarding the PRRT process
- **Goal Assessment:** Compare the number and types of interventions for activated PRRTs, ICU transfers, at staff satisfaction surveys pre- and post-implementation

- **Goal:** Implement a quality improvement initiative using the evidence-based PEWS criteria to improve recognition of deteriorating pediatric patients, allocation of PRRT resources, and pediatric staff satisfaction regarding the PRRT process
- **Goal Assessment:** Compare the number and types of interventions for activated PRRTs, ICU transfers, at staff satisfaction surveys pre- and post-implementation

PEWS (Table 2) evaluates 3 domains: behavior, cardiovascular, and respiratory; each domain ranges in point values from 0-3; a flowchart (Figure 1) has specific protocols for each score; normal VS parameters were established by age group

PEWS replaced the VS based system in June 2016; pediatric nursing staff were trained on PEWS prior to this date; pre- and post-implementation data were collected from Oct 2015 - Jun 2016 and Jul 2016 - Dec 2016, respectively

Data were collected on age, activation criteria, interventions performed, ICU transfers, code blues, potential missed opportunities, patient acuity, patient care days, and number of monthly discharges

- PEWS (Table 2) evaluates 3 domains: behavior, cardiovascular, and respiratory, each domain ranges in point values from 0-3; a flowchart (Figure 1) has specific protocols for each score; normal VS parameters were established by age group. PEWS replaced the VS based system in June 2016; pediatric nursing staff were trained on PEWS prior to this date; pre- and post-implementation data were collected from Oct 2015 - Jun 2016 and Jul 2016 - Dec 2016, respectively. Data were collected on age, activation criteria, interventions performed, ICU transfers, code blues, potential missed opportunities, patient acuity, patient care days, and number of monthly discharges. Patient acuity was estimated using nursing workload data from the Workload Management System for Nursing Internet (WMMIS). Potential missed opportunities were defined as patients meeting PRRT activation criteria that did not have a PRRT activated. PEWS was estimated in the pre-implementation group based on chart review. Pre- and post-implementation surveys were administered to all pediatric ward ICU staff regarding perceptions and confidence in the PRRT process. Survey questions were partially derived from Akre, et al¹⁰ and used a Likert scale. Surveys were analyzed using groups of favorable, neutral, and unfavorable responses, and analyzed by job title (physician, ward nursing staff, ICU staff). Surveys were excluded if unable to determine whether the staff employment start date was prior to the initial evaluation period. Categorical data were analyzed using Fisher's exact and Chi-square statistical methods; p-values <0.05 were considered statistically significant.


The island's success has been attributed to the fact that it is a

	VA Based System	PWVS
Median [IQR] Patient Age (years)	5 [2, 7, 26]	2 [1, 12]
Median [IQR] PWVS*	2 [1, 75, 5]	5 [5, 6]
# PRTTs Called	36	22
Rate of PRTTs (per 1,000 patient care days)	20.2	15.5
Pediatric Ward Code Blue Events	1	1
# Potential Missed Opportunities	28	7
Median [IQR] Patient Acuity (WME/N)	3.5 [3.3, 4.0]	3.5 [3.4, 3.5]
Mean Monthly Patient Care Days	200.3	237.2
Mean Monthly Discharges	111.5	107.5


	VA Based System	PWVS
Median [IQR] Patient Age (years)	5 [2, 7, 26]	2 [1, 12]
Median [IQR] PWVS*	2 [1, 75, 5]	5 [5, 6]
# PRTTs Called	36	22
Rate of PRTTs (per 1,000 patient care days)	20.2	15.5
Pediatric Ward Code Blue Events	1	1
# Potential Missed Opportunities	28	7
Median [IQR] Patient Acuity (WME/N)	3.5 [3.3, 4.0]	3.5 [3.4, 3.5]
Mean Monthly Patient Care Days	200.3	237.2
Mean Monthly Discharges	111.5	107.5

	0	1	2	3
Behavior	Playing Alert Appropriate of location	Sleep Family full comatose	Immobile/Incomprehensible	Lethargic Circumstantial Comprehension to part of environment
Response	Prox Capillary will 1-2 seconds	Dist Capillary will 2 seconds	Dist Capillary will 4 seconds Thumpings of 20 seconds normal rate	Dist, specific or Capillary will 4- seconds Thumpings of 30 seconds normal rate or bradycardia
Respiratory	With normal parameters No reflexes	1-15 shallow normal Adequate necessary reflexes	1-20 slower normal 40% T20 1-20 normal rate 1-15 hyperventilate	1-15 slower normal 1-15 normal rate 1-15 hyperventilate or gasping 4-1 hyperventilate

	0	1	2	3
Behavior	Playing Alert Appropriate of location	Sleep Family full comatose	Immobile/Incomprehensible	Lethargic Circumstantial Comprehension to part of environment
Response	Prox Capillary will 1-2 seconds	Dist Capillary will 2 seconds	Dist Capillary will 4 seconds Thumpings of 20 seconds normal rate	Dist, specific or Capillary will 4- seconds Thumpings of 30 seconds normal rate or bradycardia
Respiratory	With normal parameters No reflexes	1-15 shallow normal Adequate necessary reflexes	1-20 slower normal 40% T20 1-20 normal rate 1-15 hyperventilate	1-15 slower normal 1-15 normal rate 1-15 hyperventilate or gasping 4-1 hyperventilate

[illegible]

Symptom	Vira sign-based (%)	FFW5 (%)	p-value
Respiratory	~35	~85	0.001
Itch	~10	~15	
Hoarse	~10	~15	
Multiple	~35	~25	0.04
Multiple, No impairment	~10	~15	
Other	~5	~10	
Transfer to ICU	~25	~85	0.01



2015

Month	PRRF Transferred to ICU	PRRF Transferred to Higher Level of Care (NO PRRF)	PRRF Canceled	PRRF Rescheduled	PRRF Cancelled	PRRF Ratio (100 Patient Care Days)	PRRF Density (per month)
Dec	1	1	1	1	1	1.0	1.0
Jan	1	1	1	1	1	1.0	1.0
Feb	1	1	1	1	1	1.0	1.0
Mar	1	1	1	1	1	1.0	1.0
Apr	1	1	1	1	1	1.0	1.0
May	1	1	1	1	1	1.0	1.0
Jun	1	1	1	1	1	1.0	1.0
Jul	1	1	1	1	1	1.0	1.0
Aug	1	1	1	1	1	1.0	1.0
Sep	1	1	1	1	1	1.0	1.0
Oct	1	1	1	1	1	1.0	1.0
Nov	1	1	1	1	1	1.0	1.0
Dec	1	1	1	1	1	1.0	1.0

2016

Month	PRRF Transferred to ICU	PRRF Transferred to Higher Level of Care (NO PRRF)	PRRF Canceled	PRRF Rescheduled	PRRF Cancelled	PRRF Ratio (100 Patient Care Days)	PRRF Density (per month)
Oct	1	1	1	1	1	1.0	1.0
Nov	1	1	1	1	1	1.0	1.0
Dec	1	1	1	1	1	1.0	1.0
Jan	1	1	1	1	1	1.0	1.0
Feb	1	1	1	1	1	1.0	1.0
Mar	1	1	1	1	1	1.0	1.0
Apr	1	1	1	1	1	1.0	1.0
May	1	1	1	1	1	1.0	1.0
Jun	1	1	1	1	1	1.0	1.0
Jul	1	1	1	1	1	1.0	1.0
Aug	1	1	1	1	1	1.0	1.0
Sep	1	1	1	1	1	1.0	1.0
Oct	1	1	1	1	1	1.0	1.0
Nov	1	1	1	1	1	1.0	1.0
Dec	1	1	1	1	1	1.0	1.0





Figure 1 consists of five bar charts, each representing a different category. Each chart compares two groups: 'No' (blue bars) and 'Yes' (purple bars). The y-axis for all charts represents the number of responses, ranging from 0 to 30.

- Physician Demographics:**
 - MD's: No (~18), Yes (~15)
 - MD's: No (~12), Yes (~10)
 - MD's: No (~10), Yes (~8)
 - MD's: No (~8), Yes (~6)
 - MD's: No (~6), Yes (~4)
 - MD's: No (~4), Yes (~2)
 - MD's: No (~2), Yes (~1)
- Patient Demographics:**
 - No LPA/VA: No (~25), Yes (~20)
 - No LPA/VA: No (~15), Yes (~10)
 - No LPA/VA: No (~10), Yes (~5)
 - No LPA/VA: No (~5), Yes (~2)
 - No LPA/VA: No (~2), Yes (~1)
 - No LPA/VA: No (~1), Yes (~0)
 - No LPA/VA: No (~0), Yes (~0)
- Management and Prioritization for Back Patients:**
 - Agree: No (~15), Yes (~10)
 - Neutral: No (~10), Yes (~5)
 - Disagree: No (~5), Yes (~2)
 - Disagree: No (~2), Yes (~1)
 - Disagree: No (~1), Yes (~0)
 - Disagree: No (~0), Yes (~0)
 - Disagree: No (~0), Yes (~0)
- Physical Confidence in Non-Communication:**
 - Agree: No (~15), Yes (~10)
 - Neutral: No (~10), Yes (~5)
 - Disagree: No (~5), Yes (~2)
 - Disagree: No (~2), Yes (~1)
 - Disagree: No (~1), Yes (~0)
 - Disagree: No (~0), Yes (~0)
 - Disagree: No (~0), Yes (~0)
- Accuracy of RST Criteria:**
 - Agree: No (~15), Yes (~10)
 - Neutral: No (~10), Yes (~5)
 - Disagree: No (~5), Yes (~2)
 - Disagree: No (~2), Yes (~1)
 - Disagree: No (~1), Yes (~0)
 - Disagree: No (~0), Yes (~0)
 - Disagree: No (~0), Yes (~0)



- SBART data (Table 1, Figure 2, Figure 3):
 - 58 PRRTs and 2 code blue events were activated during the evaluation period
 - Post-implantation rate decreased from 20.2 to 13.5 RTs/1000 patient care days
- WMSNI data suggested that patient acuity was unchanged, although Dec 2016 was unavailable, which is typically a higher acuity month
 - Median monthly patient-care days increased from 200.33 pre-implantation to 237.17 post-implantation which confirms a higher daily ward census
 - Mean monthly hospital discharges were 111.5 pre- and 107.5 post-implantation
- During the use of the PEWS, there was an increase in clinically significant interventions ($p=0.04$), respiratory support ($p=0.001$), and ICU transfers ($p=0.01$).

PRRT Data (Table 1, Figure 2, Figure 3):

- WMSU data suggested that patient acuity was unchanged, although Dec 2016 was unavailable, which is typically a higher acuity month
- Median monthly patient-care days increased from 200.33 pre-implementation to 237.17 post-implementation which confirms a higher and higher daily census
- Mean monthly hospital discharges were 111.5 pre- and 107.5 post-implementation
- During the use of the PERSU, there was an increase in clinically significant interventions ($p=0.04$), respiratory support ($p=0.001$), and ICU transfers ($p<0.01$).

in addition to fewer potential missed opportunities

67 pre-surveys and 73 post-surveys were collected [26 (50%) pediatric physicians, 26 (84%) ward staff, and 12 (87%) ICU staff pre- and 25 (48%) physicians, 34

- Physicians reported that PEWS improved nursing communication ($p=0.02$) and more accurately identified deteriorating patients ($p=0.13$). Compared to PEWS, physicians found that the VS based system neglected signs and symptoms important to identify deteriorating patients ($p=0.0006$). Pediatric ward staff reported the PEWS improved management and prioritization of ill patients ($p=0.02$), and emphasized clinical autonomy ($p=0.07$).

PEWS implementation has been an efficient and effective means of identifying deteriorating pediatric patients on the pediatric ward. Following PEWS implementation, there was a decrease in the rate of PRRTs activated despite no change in clinical acuity and increased ward census. Use of PEWS has led to more appropriate identification of deteriorating ward patients, as evidenced by the increase in clinically significant PRRT interventions. Pediatric staff report increased confidence managing deteriorating patients and improved nursing staff clinical autonomy.

- PEWS implementation has been an efficient and effective means of identifying deteriorating pediatric patients on the pediatric ward. Following PEWS implementation, there was a decrease in the rate of PRRTs activated despite no change in clinical acuity and increased ward census. Use of PEWS has led to more appropriate identification of deteriorating ward patients, as evidenced by the increase in clinically significant PRRT interventions. Pediatric staff report increased confidence managing deteriorating patients and improved nursing staff clinical autonomy.

Continue improving PEWS system through subsequent PSDA cycles
Consider use of PEWS for pediatric patients in other areas of the hospital
Continue education and training on PEWS system for new pediatric staff

- Continue improving PEWS system through subsequent PSDA cycles
Consider use of PEWS for pediatric patients in other areas of the hospital
Continue education and training on PEWS system for new pediatric staff

Denmeijer et al. Implementation of the Pediatric Early Warning Scoring System on a Pediatric Hematology/Oncology Unit. *Journal of Pediatric Oncology Nursing* 2010;27(4):235-240.

DeWitt et al. Implementation of the Pediatric Early Warning System (BlacktipPEWS) for Nurse Identification of Deteriorating Patients. *Journal of Pediatric Nursing* 2014;19:330-340.

Alotaibi et al. Sensitivity of the Pediatric Early Warning Score to Identify Patient Deterioration. *Pediatrics* 2010; 125:e763-e769.

Good et al. Evaluating the Pediatric Early Warning Score (PEWS) system for Acutely Ill Patients in the Pediatric Emergency Department. *Society for Academic Emergency Medicine* 2014;21(11):1248-1256.

- Denmeijer et al. Implementation of the Pediatric Early Warning Scoring System on a Pediatric Hematology/Oncology Unit. *Journal of Pediatric Oncology Nursing* 2010;27(4):235-240.
- DeWitt et al. Implementation of the Pediatric Early Warning System (BlacktipPEWS) for Nurse Identification of Deteriorating Patients. *Journal of Pediatric Nursing* 2014;19:330-340.
- Alotaibi et al. Sensitivity of the Pediatric Early Warning Score to Identify Patient Deterioration. *Pediatrics* 2010; 125:e763-e769.
- Good et al. Evaluating the Pediatric Early Warning Score (PEWS) system for Acutely Ill Patients in the Pediatric Emergency Department. *Society for Academic Emergency Medicine* 2014;21(11):1248-1256.